

Canadian Bioinformatics Workshops

www.bioinformatics.ca

bioinformaticsdotca.github.io

Supported by



This page is available in the following languages:

Afrikaans বাংলাৰাশি Català Dansk Deutsch Ελληνικά English English (CA) English (GB) English (US) Esperanto
Castellano Castellano (AR) Español (CL) Castellano (CO) Español (Ecuador) Castellano (MX) Castellano (PE)
Euskara Suomi français français (CA) Galego עברית hrvatski Magyar Italiano 日本語 한국어 Macedonian Malayu
Nederlands Norsk Sesotho sa Leboa polski Português română slovenski jezik српски srpski (latinica) Sotho svenska
中文 華語 (台灣) isiZulu



Attribution-Share Alike 2.5 Canada

You are free:



to Share — to copy, distribute and transmit the work



to Remix — to adapt the work



Under the following conditions:



Attribution. You must attribute the work in the manner specified by the author or licensor (but not in any way that suggests that they endorse you or your use of the work).



Share Alike. If you alter, transform, or build upon this work, you may distribute the resulting work only under the same or similar licence to this one.

- For any reuse or distribution, you must make clear to others the licence terms of this work.
- Any of the above conditions can be waived if you get permission from the copyright holder.
- The author's moral rights are retained in this licence.



[Disclaimer](#)

Your fair dealing and other rights are in no way affected by the above.

This is a human-readable summary of the Legal Code (the full licence) available in the following languages:
[English](#) [French](#)

Gene Function Prediction

Veronique Voisin
Pathway and Network Analysis of -omics Data
July 27-29, 2020




Learning Objectives of this module

- How to use GeneMANIA
- Practical lab: website and Cytoscape app

GeneMANIA: Two types of function prediction


- “What does my gene do?” (1 gene as query)
- “Give me more genes like these” (gene list as query)

GeneMANIA in a



- Is a **functional interaction network**.
- Is a **collection of interaction networks** that are summed up into one global network. To integrate all these networks into one, there are some **weights** associated with each network that are **context dependent**.
- Is a **guilt by association/ label propagation** algorithm which finds genes and networks that are functionally associated with our **query gene list**.
- Is an interactive **web app** and a **Cytoscape app**.

Functional interaction network





Context-dependent networks

= weight of different networks used to measure node connection

Network weighting=
Automatic (pathways)

Network weighting=
Equal by network




○ query gene

Automatic: GO-based weighting assigns network weights based on how well the networks reproduce patterns of GO co-annotations (“Are genes that interact in the network more likely to have the same annotation?”),

“Function prediction: What does my gene do?”

Query : 1 gene
IPO4


Genemania web app



Function	FDR	Coverage
pore complex	1.66e-3	4 / 47
nuclear pore	1.66e-3	4 / 41

IPO4: importin 4

Import and export through the nuclear pore



Directionality of transport. The RanGTP gradient determines the vectorial transport of cargo. Translocation through the NPC is an energy-independent, reversible reaction. To achieve directionality of substrate transport, translocation is coupled to a second reaction. During import, this reaction is the dissociation of the import complex by RanGTP, and the subsequent binding of the import substrate to nuclear partners. The export process gains directionality by export complex disassembly in the cytoplasm involving GTP hydrolysis on Ran.

http://mpmp.huji.ac.il/maps/import_export.html

Default parameters

- Max resultant genes: 20
- Max resultant attributes: 10
- Network weighting
- Query-dependent weighting
- Automatically selected weighting method



○ query gene

“Give me more genes like these”

Query : 2 genes
SUZ12 and EZH1

Genemania web app

Function	FDR	Coverage
<input checked="" type="checkbox"/> ESC/E(Z) complex	2.66e-28	11 / 13
<input checked="" type="checkbox"/> PcG protein complex	3.70e-24	12 / 39
<input checked="" type="checkbox"/> histone methyltransferase complex	2.05e-21	12 / 64
<input checked="" type="checkbox"/> methyltransferase complex	1.02e-20	12 / 74
<input type="checkbox"/> NuRD complex	3.32e-6	4 / 14
<input type="checkbox"/> CHD-type complex	3.32e-6	4 / 14
<input type="checkbox"/> histone deacetylase complex	3.32e-6	5 / 42
<input type="checkbox"/> SWI/SNF superfamily-type complex	1.81e-5	5 / 63
<input type="checkbox"/> Sin3-type complex	5.27e-4	3 / 12
<input type="checkbox"/> Sin3 complex	5.27e-4	3 / 12
<input type="checkbox"/> transcriptional repressor complex	5.94e-4	4 / 54
<input type="checkbox"/> positive regulation of histone methylation	5.94e-4	3 / 13
<input type="checkbox"/> histone deacetylase activity	1.16e-3	3 / 17
<input type="checkbox"/> regulation of histone methylation	1.16e-3	3 / 17
<input type="checkbox"/> protein deacetylase activity	1.30e-3	3 / 18
<input type="checkbox"/> negative regulation of histone modification	1.98e-3	3 / 21
<input type="checkbox"/> deacetylase activity	2.83e-3	3 / 24
<input type="checkbox"/> regulation of chromosome organization	3.12e-3	4 / 90
<input type="checkbox"/> chromatin remodeling	3.23e-3	4 / 92
<input type="checkbox"/> positive regulation of histone modification	3.88e-3	3 / 28
<input type="checkbox"/> nuclear chromosome	5.50e-3	5 / 240
<input type="checkbox"/> histone binding	5.68e-3	4 / 110



[https://www.cell.com/molecular-cell/pdf/S1097-2765\(19\)30185-6.pdf](https://www.cell.com/molecular-cell/pdf/S1097-2765(19)30185-6.pdf)

 query gene

The polycomb repressive complex 2 (PRC2) is a chromatin-associated methyltransferase catalyzing mono-, di-, and trimethylation of lysine 27 on histone H3 (H3K27).


Default parameters

“Give me more genes like these”

Query : 23 genes
genes with mutations and associated with Early-Onset Alzheimer’s Disease



Function	FDR	Coverage
<input checked="" type="checkbox"/> neuron part	2.73e-4	9 / 298
<input checked="" type="checkbox"/> site of polarized growth	2.82e-4	5 / 47
<input checked="" type="checkbox"/> neuron projection	2.82e-4	8 / 233
<input type="checkbox"/> growth cone	2.82e-4	5 / 45
<input type="checkbox"/> perinuclear region of cytoplasm	4.35e-4	8 / 276
<input type="checkbox"/> Notch receptor processing	4.35e-4	4 / 21
<input type="checkbox"/> dendrite	4.39e-4	6 / 112
<input type="checkbox"/> developmental growth involved in morphogenesis	5.34e-4	5 / 61
<input type="checkbox"/> neuronal cell body	6.06e-4	5 / 64
<input type="checkbox"/> cell body	7.24e-4	5 / 69
<input type="checkbox"/> receptor complex	7.24e-4	7 / 213
<input type="checkbox"/> membrane protein ectodomain proteolysis	1.29e-3	4 / 33
<input checked="" type="checkbox"/> beta-amyloid metabolic process	1.64e-3	3 / 10
<input type="checkbox"/> dendritic shaft	1.64e-3	3 / 10
<input type="checkbox"/> neuron death	2.49e-3	5 / 94
<input type="checkbox"/> membrane protein proteolysis	2.69e-3	4 / 43
<input type="checkbox"/> axon	2.69e-3	5 / 97
<input type="checkbox"/> developmental growth	4.93e-3	5 / 112
<input type="checkbox"/> negative regulation of multicellular organismal process	1.08e-2	6 / 230
<input type="checkbox"/> regulation of anatomical structure size	1.90e-2	5 / 151
<input type="checkbox"/> neuron apoptotic process	2.63e-2	4 / 80
<input checked="" type="checkbox"/> autophagy	5.47e-2	4 / 98



○ query gene

Default parameters

Max resultant genes: 20

Max resultant attributes: 10

Network weighting


Query-dependent weighting

Automatically selected weighting method

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6471359/>

“Give me more genes like these”

Query : 43 genes
cholesterol metabolism related genes





Larger nodes: query genes
smaller white nodes: related genes
Red: up-regulation
Blue: down-regulation
2 time points: 3h and 21h (node center and node border)

Model of Atherosclerosis

Quiescent human coronary artery SMCs were treated with minimally-oxidized LDL (moxLDL), for 3 hours and 21 hours, respectively.

“from EnrichmentMap to GeneMANIA”

Query : 502 genes
nc RNA metabolic process




- Right click on pathway of interest
- Select 'Apps'
- Select 'EnrichmentMap – Show in GeneMANIA'

Node size and color: relative to gene expression

“Advanced topics”

- Possibility to add your model organism if you can construct or have access to interaction networks for it.
- Possibility to use GeneMANIA as an example to assemble different interaction networks that you might have for your research question.

Example: netDx is a **patient classifier** algorithm based on the GeneMANIA algorithm



Acknowledgment

Quaid Morris for his slides and lecture:

<https://www.youtube.com/watch?v=2KrUq9ad2xc&list=PL3izGL6oi0S9amh6Cza1bUo4ICcvZlWF0&index=6>

GeneMANIA developer team: Christian Lopez, Max Franz, Ruth Isserlin, Quaid Morris , Gary Bader

We are on a Coffee Break & Networking Session

compute | calcul
canada | canada

Workshop Sponsors:



Canadian Centre for
Computational
Genomics

MiCM McGill initiative in
Computational Medicine